


8. Dispersion processes



This section looks at the longitudinal spreading of clouds through dispersion, a process distinct from diffusion. Dispersion is the spreading of a cloud by any process that causes different chemical particles to move at different velocities. The non-uniformity of particle velocity allows some particles to move quickly while others are held back, and can be caused by boundaries (which create shear in the flow), dead zones and absorption reactions (for example, to a porous medium). In environmental systems, dispersion is generally far more important than diffusion in the lengthening of chemical clouds. In this chapter, it is shown that dispersing clouds eventually reach a Fickian limit, at which point they have a Gaussian concentration profile in the longitudinal direction. It is also shown that longitudinal dispersion is maximized when lateral diffusion is small.

The example problems ask the user to make qualitative statements about the dispersion expected in various flows and to predict the shape of a dispersing cloud.

8. Shear Dispersion.

The transport models and concentration field solutions developed in previous sections assume that currents are spatially uniform, *i.e.* $u \neq f(x,y,z)$. However, spatial gradients of velocity, known as shear, must exist close to a boundary because the no-slip condition requires that the relative velocity of the fluid be zero at the boundary. When shear is present, different parts of a tracer cloud will advect at different speeds. This 'differential advection' stretches the tracer cloud, causing it to spread longitudinally more quickly than a cloud released into a uniform current (Figure 1). In addition, differential advection creates lateral gradients in concentration ($\partial C/\partial y$ in Figure 1) which, when acted upon by lateral diffusion, accelerates the dilution of the cloud.




Figure 1. Top-view of two channels. In the top channel the side-wall boundary condition allows slip, and the velocity profile is uniform across y . In the bottom channel there is a no-slip condition at the side-wall boundary, creating lateral shear, $\partial u/\partial y$. Both channels have the same cross-sectional mean velocity, \bar{u} . Tracer is released at $t = 0$, $x = 0$. The spatial distribution of each cloud is shown above at time t_1 and t_2 . Note, the center of mass for both clouds advects at speed \bar{u} . Velocity shear in the bottom channel stretches the tracer cloud, and as a result this cloud spreads longitudinally more rapidly than the cloud in the uniform channel.

The distortion observed in the bottom channel of Figure 1 can be predicted from the full transport equation,

$$\frac{\partial C}{\partial t} + u \frac{\partial C}{\partial x} + v \frac{\partial C}{\partial y} + w \frac{\partial C}{\partial z} = \frac{\partial}{\partial x} D_x \frac{\partial C}{\partial x} + \frac{\partial}{\partial y} D_y \frac{\partial C}{\partial y} + \frac{\partial}{\partial z} D_z \frac{\partial C}{\partial z}, \quad (1)$$

if we include the spatial variation in velocity. However, with $u = f(x,y,z)$, it is difficult to find simple analytical solutions to (1). In many instances it is sufficient to integrate the domain over y and z , and consider only transport in the stream-wise direction. After this integration the combined impact of differential advection and lateral diffusion is

expressed as an additional process creating longitudinal spreading, which we term Shear Dispersion. It is also known as Taylor Dispersion, in honor of G. I. Taylor, who first described this process in 1953. Under limiting conditions, to be discussed shortly, Shear Dispersion behaves as a Fickian diffusion process. Thus, we represent its impact in the cross-sectionally averaged (one-dimensional) transport equation with an additional coefficient of dispersion, K_x . Integrating (1) over y and z yields the one-dimensional equation,

$$\frac{\partial \langle C \rangle}{\partial t} + \langle u \rangle \frac{\partial \langle C \rangle}{\partial x} = D_x \frac{\partial^2 \langle C \rangle}{\partial x^2} + K_x \frac{\partial^2 \langle C \rangle}{\partial x^2}, \quad (2)$$

in which brackets represent cross-sectional averages, and for simplicity we have assumed homogeneous diffusivity, $D_x \neq f(x)$. In general $D_x \ll K_x$, such that the former may be neglected, leaving us with the one-dimensional advection-dispersion equation.

$$\frac{\partial \langle C \rangle}{\partial t} + \langle u \rangle \frac{\partial \langle C \rangle}{\partial x} = K_x \frac{\partial^2 \langle C \rangle}{\partial x^2}. \quad (3)$$

Furthermore, in the limit of Fickian behavior, we describe the evolution of the cross-sectionally averaged cloud using the same statistical description developed for diffusion in previous chapters. Specifically, the longitudinal length-scale for the cloud is σ_x , where

$$\sigma_x = \sqrt{2 K_x t}. \quad (4)$$

From (4) we can estimate K_x in the field by observing the evolution of variance for a tracer cloud. Specifically, observing the variance over time,

$$K_x = \frac{1}{2} \frac{\partial \sigma_x^2}{\partial t}. \quad (5)$$

We can find an analytical expression for K_x by explicitly carrying out the integration that leads to (3), utilizing scaling arguments to simplify the mathematics along the way. It is important to understand the steps of this derivation, to appreciate under what conditions the approximate equation (3) may hold. For simplicity we will consider a very wide channel for which the shear in the lateral, $\partial u / \partial y$, is so small compared to the shear in the vertical, $\partial u / \partial z$, that the lateral shear is neglected at the outset. This allows us to assume a two-dimensional channel with $\partial / \partial y = 0$. In addition, we assume that the flow is strictly in the x -direction ($w = v = 0$), and uniform ($\partial u / \partial x = 0$). With these assumptions, (1) reduces to

$$\frac{\partial C}{\partial t} + u \frac{\partial C}{\partial x} = \frac{\partial}{\partial x} D_x \frac{\partial C}{\partial x} + \frac{\partial}{\partial z} D_z \frac{\partial C}{\partial z}. \quad (6)$$

Next, we decompose u and C into depth-averaged values denoted by an over-bar, e.g. $\bar{u} = (1/h) \int_0^h u dz$. Deviations from the depth-average will be denoted by a prime.

$$u(z) = \bar{u} + u'(z) \tag{7}$$

$$c(x,z) = \bar{c}(x) + c'(x, z)$$

We put the decomposed variables into (6), and note that by definition $\partial c/\partial z = 0$.

$$\frac{\partial}{\partial t}(\bar{c} + c') + (\bar{u} + u') \frac{\partial(\bar{c} + c')}{\partial x} = \frac{\partial}{\partial x} D_x \frac{\partial(\bar{c} + c')}{\partial x} + \frac{\partial}{\partial z} D_z \frac{\partial(c')}{\partial z}. \tag{8}$$

We take the depth-average of each term, noting that by definition $\overline{c'} = 0$ and $\overline{u'} = 0$.

$$\frac{\partial \bar{c}}{\partial t} + \bar{u} \frac{\partial \bar{c}}{\partial x} + \overline{u' \frac{\partial c'}{\partial x}} = \frac{\partial}{\partial x} D_x \frac{\partial \bar{c}}{\partial x} \tag{9}$$

Relative to our original equation, (6), we have eliminated the vertical gradient terms by integrating over the vertical domain. In addition, a new term has appeared, $\overline{u' \partial c' / \partial x}$, that represents the flux associated with the correlation between the fluctuations in velocity, u' , and concentration c' , relative to their depth-averaged values. With the next few steps we will find a solution for c' , so that we might evaluate this new flux term. As both (8) and (9) must both hold true, we may subtract (9) from (8) to yield.

$$\frac{\partial c'}{\partial t} + u' \frac{\partial \bar{c}}{\partial x} + u' \frac{\partial c'}{\partial x} - \overline{u' \frac{\partial c'}{\partial x}} = \frac{\partial}{\partial z} D_z \frac{\partial c'}{\partial z} \tag{10}$$

Now we place an important condition on the magnitude of the spatial fluctuation c' . Specifically, we assume that $c' \ll \bar{c}$. If this is true, then the magnitude of both the third and fourth terms in (10) must be small compared to the second term, and we may drop the small terms, reducing (10) to

$$\frac{\partial c'}{\partial t} + \underbrace{u' \frac{\partial \bar{c}}{\partial x}}_{\text{differential advection}} = \underbrace{\frac{\partial}{\partial z} D_z \frac{\partial c'}{\partial z}}_{\text{vertical diffusion}} \tag{11}$$

This equation describes how vertical fluctuations in concentration, c' , are created by differential advection and destroyed by vertical diffusion. Initially, when $\partial \bar{c} / \partial x$ is large, differential advection dominates, creating c' faster than vertical diffusion can erase it. But, eventually, $\partial \bar{c} / \partial x$ becomes small enough that the rate at which c' is created by differential advection is balanced by the rate at which it is destroyed by vertical diffusion. When these two processes are in balance, $\partial c' / \partial t$ goes to zero and (11) becomes.

$$u' \frac{\partial \bar{c}}{\partial x} = \frac{\partial}{\partial z} D_z \frac{\partial c'}{\partial z}. \tag{12}$$

We can solve (12) with a no-flux boundary condition at the bed and water surface, *i.e.* $\partial c'/\partial z = 0$ at $z = 0$ and h . Recall that by definition, $\bar{c} \neq f(z)$. Then,

$$c'(z) = \frac{\partial \bar{c}}{\partial x} \int_0^z \frac{1}{D_z} \int_0^z u' dz dz + c'(z=0). \quad (13)$$

Because the diffusivity, D_z , is the sum of turbulent and molecular diffusion, it may be a function of vertical position, and so remains inside the integral.




Figure 2. Consider a tracer cloud whose leading edge is uniform at time t . Differential advection distorts the cloud giving it a new shape by time $t+dt$. The spatial deviation in velocity, u' , creates the concentration perturbation seen at $t+dt$. Specifically, from the first two terms in (11), $c' = -(\partial \bar{C}/\partial x) u' dt$.

Now we return to (9) to evaluate the new term, $\overline{u' \partial c'/\partial x}$. Because we assume $u \neq f(x)$, then $u' \neq f(x)$, and we can write the new term as $\overline{\partial u' c'/\partial x}$. We evaluate the correlation $\overline{u' c'}$ with (13), noting that $\overline{u' c'(z=0)} = \overline{u' c'(z=0)} = 0$, because $\overline{u'} = 0$.

$$\overline{u' c'} = \frac{1}{h} \int_0^h u' \frac{\partial \bar{c}}{\partial x} \int_0^z \frac{1}{D_z} \int_0^z u' dz dz dz = \frac{1}{h} \frac{\partial \bar{c}}{\partial x} \int_0^h u' \int_0^z \frac{1}{D_z} \int_0^z u' dz dz dz. \quad (14)$$

This correlation is the advective flux associated with spatial fluctuations in the velocity field. This flux is proportional to the concentration gradient, and so behaves as a Fickian diffusion. Thus, we choose to model this flux using a transport coefficient, just as we did for turbulent diffusion in the previous chapter. The transport coefficient is called the longitudinal dispersion coefficient, K_x . The negative sign indicates that the flux is counter-gradient.

$$\overline{u' c'} = -K_x \frac{\partial \bar{c}}{\partial x}, \quad (15)$$

where

The second expression in (16) is valid when $D_z \neq f(z)$. This expression describes the contribution of vertical shear to longitudinal dispersion. A similar term describes the contribution of lateral shear.

Now we can write the new term in (9) as,

$$\overline{u' \frac{\partial c'}{\partial x}} = \frac{\partial}{\partial x} \overline{u' c'} = -K_x \frac{\partial^2 \bar{c}}{\partial x^2} \quad (17)$$

Using (17) in (9), we arrive at the one-dimensional equation,

$$\frac{\partial \bar{c}}{\partial t} + \bar{u} \frac{\partial \bar{c}}{\partial x} = \frac{\partial}{\partial x} D_x \frac{\partial \bar{c}}{\partial x} + K_x \frac{\partial^2 \bar{c}}{\partial x^2}, \quad (18)$$

where D_x in (18) and D_z in (16) represent the sum of molecular and turbulent diffusion, which for practical purposes is just turbulent diffusion. Elder evaluated (16) for a logarithmic profile with $D_z = f(z)$ and found

$$K_x = 5.93u_*h. \quad (19)$$

Note that (19) only accounts for velocity shear over depth ($\partial u/\partial z$) and not for lateral velocity shear ($\partial u/\partial y$). This is an acceptable approximation for wide, straight channels. Comparing (19) with the empirical estimate for longitudinal diffusion, $D_x = 0.45u_*h$, we see that K_x is an order of magnitude larger. Because of this, the diffusion term is often neglected in (18), and we have returned to equation (3), repeated here for convenience.

$$\frac{\partial \bar{c}}{\partial t} + \bar{u} \frac{\partial \bar{c}}{\partial x} = K_x \frac{\partial^2 \bar{c}}{\partial x^2} \quad (20)$$

Solutions to this one-dimensional equation, with D_x replaced by K_x , have already been described for instantaneous and continuous point releases. These solutions can now be used for systems with shear-dispersion dominating longitudinal diffusion (as in 20). As one might expect, (16) shows that the coefficient of shear dispersion increases with the magnitude of the shear, expressed as the magnitude of the spatial deviations u' . The dependence $K_x \sim h$ is less straight-forward. However, considering that each integral is proportional to h , and that $D_z \sim h$, (16) yields $K_x \sim h$, consistent with (19). Finally, (16) indicates that K_x is inversely proportional to the cross-stream diffusivity, D_z (i.e. K_x is greatest when D_z is smallest!). To understand this somewhat unintuitive dependence, remember that shear dispersion arises because velocities on adjacent streamlines are different so particles on different streamlines advect at different speeds and thus become separated (dispersed) longitudinally. This is the differential advection component of shear dispersion. At the same time, each particle is mixed over depth (or width), experiencing the velocity of each streamline that it passes for some fraction of time. If

the cross-channel diffusion is high, the particles sample all the velocities in the profile so rapidly that they effectively travel at the mean channel velocity. If they all effectively travel at the mean velocity, they do not disperse (separate) in the longitudinal direction, and shear dispersion will be zero, even though shear is not zero. As the cross-stream diffusion is reduced, particles can accumulate more differential advection as they remain on individual streamlines for a longer period.

Now it is important to recall the assumptions that allowed us to reach the advection-dispersion equation (20). Early on we made the assumption that $c' \ll \bar{c}$. In addition, for (12) to be valid, we specifically required that the rate at which c' is created exactly balance the rate at which c' is destroyed, so that a nearly uniform concentration profile exists in the cross-stream direction. Note that a nearly uniform profile would automatically satisfy $c' \ll \bar{c}$, such that the two assumptions are closely linked. To understand when these assumptions are valid, we consider the evolution of a tracer cloud from its initial release, as shown in Figure 3. The tracer is introduced as a line distributed over the full depth at time $t = 0$. Differential advection, u' , creates perturbations in the concentration profile, c' , at the rate $\partial c' / \partial t = u' \partial \bar{c} / \partial x$. When the cloud is first introduced, the mean concentration gradients are very large, such that the creation of c' is very rapid and pronounced. This is reflected in the geometric distortion of the cloud shape (**t1**, Fig. 3). The concentration perturbations are dissipated by lateral diffusion, which acts at the rate, $\partial c' / \partial t = D_z \partial^2 c' / \partial z^2$. As the cloud spreads, the longitudinal gradient of mean concentration, $\partial \bar{c} / \partial x$, decreases and the rate at which concentration perturbations are created slows. Eventually, the rate at which the perturbations are created by differential advection comes into balance with the rate at which they are destroyed by diffusion such that (12) is valid. From tracer studies, this is observed to occur at time scale

$$T_{\text{Dispersion}} = 0.4 L_z^2 / D_z, \quad (21)$$

after the tracer is released, where L_z is the length-scale of the channel in the cross-stream direction (width or depth). For vertical shear, $L_z = h$, the water depth. After this time (e.g. **t2** in Figure 3) the concentration is close to uniform over depth, such that $c' \ll \bar{c}$. The process of shear-dispersion has reached its Fickian limit and the longitudinal distribution of the depth-averaged concentration is Gaussian. Note that this time scale is inversely proportional to cross-stream diffusion, D_z . And recall from (16) that K_x is also inversely proportional to D_z . This means that the conditions (D_z) which yield the greatest K_x will also require the longest time (longest distance from source) for shear-dispersion to take hold and for (20) to apply.




Figure 3. Evolution of tracer cloud released at $t = 0$ as a line distributed over depth. Initially (t_1) differential advection dominates, visibly distorting the geometric shape of the cloud. At this time (t_1) the longitudinal distribution of the depth-averaged concentration, \bar{c} , is skewed. Eventually differential advection and diffusion balance and the cloud approaches uniform concentration over depth. After this time (t_2) the longitudinal distribution of the depth-averaged concentration is Gaussian and the dispersion process has reached its Fickian limit.

EXERCISE WITH THE SOLUTION

Describe the following processes: diffusion, advection and dispersion

SOLUTION

Diffusion is a specific term referring to the net flux resulting from zero-mean, random motions. The random motions may be Brownian motion – resulting in molecular diffusion – or may be due to turbulent eddies. Strictly, the size of the eddies must be small compared to the patch of particles in order for impact of the eddies on the patch to be truly random, and thus the net flux to be purely diffusive.

Advection is the transport of particles (concentration) by the mean current.

Dispersion is a broad term used to describe all processes, except diffusion, that act to disperse a patch, and thus diminish concentration. Typically, dispersion coefficients reflect a combination of advection and diffusion processes that are difficult to model separately. For example, the coupled effect of differential longitudinal advection and cross-channel diffusion creates shear dispersion. The existence of multiple discrete flow paths, e.g. through a porous medium, also creates a dispersion of particles, because the time to traverse each path differs with its geometry and the degree of bending/turning – often called the tortuosity. Thus, many particles released together but which take different pore pathways will get separated (dispersed) in the longitudinal direction. Finally, the lateral diffusion of particles into a region of zero velocity (a pore space with no outlet, a side-pool in a river, a wake behind an obstruction) will create longitudinal dispersion. The particles that enter the zones of zero velocity get held back relative to the particles that do not get trap. When the particles are eventually released from the ‘dead-zone’, they are separated in space (dispersed) from the rest of the cloud.