


Discrete Mathematics
Lecture 9
Relations and Functions

Lecturer: Kahenya, N.P




Introduction to lecture 9



Relations and functions are fundamental concepts in mathematics and computer science and in particular graph theory.



This lecture forms a foundation for the lecture on trees and graph theory in general.



Relations can be used to understand and solve problems in communications, networks, relationships between members of a social group among others.

References

These lecture notes have been derived from the following sources (Lipschutz & Lipson, 2007; Rosen, 2012).



Intended Learning Outcomes

Be	At the end of this lecture, you will be able to;
Explain	Explain relations and functions.
Apply	Apply the concepts in solving problems involving relations and functions.

Introduction to Relations

Relations are structures that are used to represent the relationships between elements of a given set. Relations can be used to solve problems such as;

Determining which pairs of towns that connected by roads or trains.

Determining how a set of computers are networked.

Determine the best way to store information in a computer database.

Binary relations

□ Product sets are examples of binary relations where it is used to express relationship between elements of two sets.

□ For example, let A and B be any two sets.

The set of all ordered pairs of (a, b) where $a \in A$ and $b \in B$ is called the product or Cartesian product of A and B i.e.

$$A \times B = \{(a, b) | a \in A, b \in B\}$$

□ For example, $\mathbb{R} \times \mathbb{R} = \mathbb{R}^2$ is the set of ordered pairs of real numbers.



Example 1

Let $A = \{2,3\}$ and $B = \{a, b, c, d\}$ then;

$$A \times B = \{(2, a), (2, b), (2, c), (2, d), (3, a), (3, b), (3, c), (3, d)\}$$

$$B \times A = \{(a, 2), (a, 3), (b, 2), (b, 3), (c, 2), (c, 3), (d, 2), (d, 3)\}$$

$$A \times A = \{(2,2), (2,3), (3,2), (3,3)\}$$

Note that;

(i) $A \times B \neq B \times A$

(ii) The order in which the elements appear matters.

(iii) $n(A \times B) = n(A) \cdot n(B)$ where $n(A)$ is the cardinality of set A.

Definition 1

Let A and B be sets. A binary relation or simply a relation from A to B is a subset of $A \times B$.

That is, for each pair $a \in A$ and $b \in B$ exactly one of the following is true;

(i) $(a, b) \in R$ i.e. 'a is R-related to b' written $a R b$.

(ii) $(a, b) \notin R$ i.e., 'a is not R-related to b' written $a \not R b$

Remark 1:

If R is a relation from a set A to itself, that is if R is a subset of $A^2 = A \times A$ then we say that R is a relation on A .



Example 1

Let $A = \{1,2\}$ and $B = \{x,y\}$ and

$R = \{(1,x), (1,y)\}$

then R is a relation from A to B since R is a subset of $A \times B$.

Hence, we have $1 R x, 1 R y, 2 \not R x, 2 \not R y$.

Note that the domain of R is $\{1\}$ and the range of R is $\{x,y\}$.

Example 2

Let $A = \{\text{students in a class}\} = \{\text{Oti, Onyi, Kama, Chris, Njeri}\}$ and B be the set of courses i.e.

$B = \{\text{History, Maths, English}\}.$

Let R be the relation that consists of the pairs (a,b) where a is a student enrolled in course b .

For instance, if Oti and Njeri are enrolled in maths then we have (Oti, Maths) and (Njeri, Maths) belong to R .

Examples of Relations

- 1) Set inclusion \subseteq is a relation on any collections of sets e.g., $A \subseteq B$ or $A \not\subseteq B$.
- 2) In the set of integers, we have such relations as $m|n$ i.e., m divides n e.g., $4|12$, $5|25$ but $4 \nmid 17$.
- 3) Functions are relations. Where a function says, f from set X to a set Y assigns exactly one element of Y to each element of X . The graph of f is a set of ordered pairs (x, y) so that $y = f(x)$ which is a subset of $X \times Y$.

Inverse relation

Let R be any relation from set A to set B .

The inverse of R denoted R^{-1} is the relation from B to A which consists of ordered pairs which when reversed belong to R i.e.,

$$R^{-1} = \{(y, x) \mid (x, y) \in R\}.$$

Example 1: Let set $A = \{1, 2\}$ and $B = \{a, b\}$ then the inverse of $R = \{(1, a), (1, b), (2, a)\}$ is $R^{-1} = \{(a, 1), (b, 1), (a, 2)\}$.



Example 2

Consider a set A with n elements. A relation on the set A is a subset of the $A \times A$.

The set $A \times A$ has n^2 elements.

The set A with n elements has 2^n subsets, and therefore we can say that $A \times A$ has 2^{n^2} subsets.

For instance, the set $A = \{3, d, f, g\}$ has $2^{4^2} = 65536$ relations.

Representation of Relations


a) Matrix of relation

This can be done in a matrix form where we use 1 or 0 if there is a relation or no relation respectively.


For example, let sets $A = \{1,2,3\}$ and $B = \{a, b, c\}$ and $R = \{(1, b), (1, c), (2, c), (3, a)\}$.

Then the matrix of relation is $M_R = \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$

	a	b	c
1	0	1	1
2	0	0	1
3	1	0	0



In general, if $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$ are finite sets containing m and n elements, respectively. R is a relation from A to B then R can be represented by the $m \times n$ matrix $M_R = [m_{ij}]$ i.e.


$$M_R = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$



The matrix M_R is called the matrix of R .

Example 1

Let $M_R = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}$ and let sets $A = \{a, b, c\}$ and $B = \{w, x, y, z\}$.


Then by definition $(a_i, b_j) \in R$ iff $m_{ij} = 1$ thus;

$R = \{(a, w), (a, z), (b, x), (b, y), (c, x), (c, z)\}$.




Directed graphs/digraphs of R

Relations can be represented using directed graphs, where the elements represent the vertices or nodes of the digraph and the relation between vertices is represented by a directed edge.



Example 1

Let R be a relation on the set $A = \{1,2,3,4\}$ where $R = \{(1,2), (2,2), (2,4), (3,4), (4,1), (4,3)\}$ then its directed graph is;



Example 2 box


Let $A = \{1,2,3,4\}$ and $R = \{(1,1), (1,3), (1,4), (2,3), (3,1), (3,4), (4,4), (2,4)\}$.

Then the diagraph of the relation is;



Example 3

Find the relation determined by digraph below;



Definition 1

The in-degree of a vertex is the number of edges terminating at the vertex or node or junction, while the out-degree of a vertex is the number of edges leaving a vertex.



Example 3

Let set $A = \{1,2,3,4\}$ and let the relation on A represented by the

$$\text{matrix, } M_R = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \end{pmatrix}$$

Construct the digraph of R and list in-degree and out-degrees of all vertices.



- **Graphs of functions**

Functions are relations and hence can be represented by graphing.


- **Arrow diagram**

Relations can be represented in an arrow diagram.

For example, let sets $A = \{1,2,3\}$ and $B = \{a, b, c\}$ and

$R = \{(1, b), (1, c), (2, c), (3, a)\}$.

Composition of relations

A network diagram on a blue background. It consists of several white pushpins connected by black lines, forming a complex web. One pushpin in the center is red, standing out from the others.

Definition 1: Let A , B , and C be any non-empty sets. Let R and S represent the relations from A to B , and B to C respectively. In other words, the relation R is a subset of $A \times B$ while the relation S is a subset of $B \times C$.

Then the relation from A to C denoted $S \circ R$, the composition of S and R , is defined as;

$$a(S \circ R)c$$

If for some $b \in B$ then aRb and bSc , that is,

$$S \circ R = \{(a, c) | \exists b \in B, (a, b) \in R, (b, c) \in S\}$$

Example 1 box

Consider the sets $A = \{1,2,3\}$, $B = \{a, b, c\}$ and $C = \{\alpha, \beta, \gamma\}$ and the relations $R = \{(1, b), (1, c), (3, a), (2, c)\}$ and $S = \{(a, \alpha), (b, \beta), (c, \alpha), (c, \beta)\}$. Determine $S \circ R$.

From the arrow diagram we get; $S \circ R = \{(1, \alpha), (1, \beta), (2, \alpha), (2, \beta), (3, \alpha)\}$



Alternatively

Alternatively, consider the matrix of relations R and S i.e. $R = \{(1, b), (1, c), (3, a), (2, c)\}$ and $S = \{(a, \alpha), (b, \beta), (c, \alpha), (c, \beta)\}$.

$$M_R M_S = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix} = \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} \begin{bmatrix} 1 & 2 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

The 2 implies that there are two ways from 1 to β . The non-zero entries indicate the relation $S \circ R$ exists.

Types of relations

REFLEXIVE


SYMMETRIC

Asymmetric

Antisymmetric

TRANSITIVE

Equivalence



Reflexive

A relation R on a set A is called reflexive if $(a, a) \in R$ for every element $a \in A$.

That is R is reflexive if every element $a \in A$ is related to itself, otherwise it is irreflexive.

Example 1: Consider the set $A = \{a, b, c\}$ and the relations R on the set A namely;

$$R_1 = \{(a, a), (a, b), (a, c), (c, c)\}$$

$$R_2 = \{(a, b), (b, b), (c, c), (a, a), (a, c)\}$$

$$R_3 = \{(a, a), (b, b), (c, a), (c, b)\}$$

Only R_2 is reflexive since we have $(a, a) \in R$, $(b, b) \in R$, $(c, c) \in R$.

Note that the universal relation $A \times A$ is also reflexive.

Symmetric

A relation R on a set A is symmetric if whenever $a R b$ then $b R a$.

A relation R on a set A is asymmetric if whenever $a R b$ then $b R a$ i.e., $(a, b) \in R$ then $(b, a) \notin R$.

A relation R on a set A is antisymmetric if whenever $a R b$ and $b R a$ then $a = b$. Also, R is still antisymmetric if whenever $a \neq b$ then $(a, b) \notin R$ or $(b, a) \notin R$.

Remarks

Symmetric is not the opposite of antisymmetric.

A relation can have both of these properties or may lack.

A relation can be both symmetric and antisymmetric if it contains some pair of the form (a, b) , with $a \neq b$

Example 1

Let $A = \mathbb{Z} = \{\text{set of integers}\}$ and let $R = \{(a, b) \in A \times A \mid a < b\}$ where R is the relation less than. Is R symmetric, asymmetric, or antisymmetric?

Solution:

Symmetry: If $a < b$ then it is not true that $b < a$. Therefore, R is not symmetric.

Asymmetry: If $a < b$ then $b \not< a$ so R is asymmetric.

Antisymmetric: If $a \neq b$ then either $a < b$ or $b < a$. Therefore,

R is antisymmetric.

Transitive

A relation R on a set A is called transitive if whenever $(a, b) \in R$ and $(b, c) \in R$ then $(a, c) \in R$ for all $a, b, c \in A$ i.e. If $a R b$ and $b R c$ then $a R c$.

Example 1: Is the 'divides' relation on set of positive integers transitive?

Solution: Let a divides b and b divides c , does a divides c ?

If a divides b then there exists an m such that $b = am \cdots$ (i)

Also, if b divides c then there exists an n such that $c = bn \cdots$ (ii)


From (i) and (ii) we have; $c = bn = (am)n = a(mn) \Rightarrow a|c$. Hence 'divides' relation is transitive.

Functions

- Let X and Y be sets. A function from X to Y is a rule that assigns to each element of X exactly one element of Y .
- A function is generally denoted with f , g , h etc.
- If $f: X \rightarrow Y$ then X is called the Domain of f and Y is the codomain.
- If $f: X \rightarrow Y$ the set of images $\{y \in Y: y = f(x) \text{ for some } x \in X\}$ is called the range of f .
- Note that range is a subset of codomain.

Invertible functions

- A function f from set A to B is one-to-one if no two distinct elements of the domain have the same image i.e. If $f(a) = f(b)$ then $a = b$.
- A function f from set A to set B is onto if $\forall b \in B, \exists a \in A$ such that $f(a) = b$
- **Theorem 2:** A function f has an inverse iff f is onto and 1-1.
- A function $f: X \rightarrow Y$ is said to be invertible if its inverse relation f^{-1} is also a function.
- Note that a function is not necessarily invertible.



Theorem 1

Let $f: X \rightarrow Y$ be a function

(i) Then f^{-1} is a function from B to A iff f is 1-1.

(ii) If f^{-1} is a function, then the function f^{-1} is also 1-1.

(iii) f^{-1} is everywhere defined iff f is onto.

(iv) f^{-1} is onto iff f is everywhere defined.



Composite function

Consider the functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ i.e., where the codomain of f is domain of g .

We define the function from X to Z as the composition of f and g written $f \circ g$ as follows

$$(g \circ f)(x) = g(f(x))$$



Inverse of a function

Let f be 1 – 1 function, then g is the inverse of f denoted f^{-1} if;

$$(f \circ g)(x) = x, \forall x \in D(g)$$

$$(g \circ f)(x) = x, \forall x \in D(f)$$

Determining the inverse of $y = f(x)$

Steps

- ❑ Check if f is 1 – 1
- ❑ If yes, then solve for x i.e., let $x = f^{-1}(y)$.
- ❑ Exchange x and y to get $y = f^{-1}(x)$
- ❑ Check that $(f \circ f^{-1})(x) = x = (f^{-1} \circ f)$

Example 1: Determine whether $f(x) = 5x$ is a 1 – 1 function

Solution: Suppose $f(x) = \alpha \Rightarrow \alpha = 5x \therefore x = \frac{\alpha}{5} \Rightarrow f(x)$ is 1 – 1.

Example 2: Determine if $f(x) = x^2$ is a one-to-one function.

Solution: Let $f(x) = \alpha \therefore \alpha = x^2 \Rightarrow x = \pm\sqrt{\alpha}$. Therefore $f(x)$ is not one-to-one (since we are getting two values of x).

Example 3 dead log

If f be defined by $f(x) = x^3 - 2$ and $g(x) = \sqrt[3]{(x+2)}$. Determine if g is the inverse of f .

Solution:

Step 1: Determine if f is 1-1. Let $f(x) = \alpha \Rightarrow x^3 - 2 = \alpha \therefore x = \sqrt[3]{(x+2)}$ i.e. f is 1-1.

Step 2: By definition if g is the inverse of f then $(f \circ g)(x) = (g \circ f)(x) = x$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt[3]{(x+2)}) = (\sqrt[3]{(x+2)})^3 - 2 = x$$

$$(g \circ f)(x) = g(f(x)) = g(x^3 - 2) = \sqrt[3]{(x^3 - 2) + 2} = x$$
$$\Rightarrow g = f^{-1}$$

Applications of Relations

- ❑ It is used in relational data model, where a database consists of records that are n-tuples and are made up of fields.
- ❑ Tables are used to show relations in a database.
- ❑ The database query language SQL (Structured Query Language) is used to carry out the operations of relations in a database.

References



Lipschutz, S., & Lipson, M. (2007). *Discrete Mathematics*. McGraw-Hill.



Rosen, K. (2012). *Discrete mathematics and its application* (7th ed.). McGraw-Hill.