


1. Dialysis is common laboratory procedure used to remove low molecular weight solutes from protein solutions. Example situations include: removal of free, unreacted label after fluorescently labeling a protein; and removal of urea from a denatured protein solution to allow refolding of the protein. The protein solution is sealed inside a dialysis bag or tubing and placed in container with a large excess of pure water, as shown in the figure. The system is then allowed to come to equilibrium. For this problem, consider an idealized system where the only solute in the dialysis bag is the low molecular weight solute (i.e., no protein is actually present.)
- Explain why the combined system of the protein solution (system “A”) and the water (system “B”) may be considered to be an isolated system.
 - Use a lattice model, with each individual water molecule occupying a single site in the lattice, to show that the entropy of System B is zero at the start of the experiment. Designate the number of water molecules N_{WB} and the number of total lattice sites M_B .
 - Use a lattice model to express the entropy of system A at the start of the experiment in terms of the number of water molecules in system A, N_{WA} , the number of solute particles, N_s , and the total number of lattice sites in system A, M_A . For your final expression, use an appropriate approximation to eliminate factorial expressions. You may assume that the water molecules and solute molecules are comparable in size.
 - Now write an expression for the entropy of the combined system under the conditions where water and solute can freely pass through the dialysis membrane. For every solute molecule that crosses over to system B, a water molecule must cross over to system A. (You may want to keep track of the number of solute molecules by noting that $N_s = N_{sA} + N_{sB}$ and the number of water molecules by noting that $N_{WA} + N_{WB} = N_W$).
 - Show that you can write the entropy expression in terms of one system variable (i.e., one parameter that changes value during the progress of the experiment), and briefly describe how to determine the equilibrium condition in terms of that variable and constants in the system. [You do not need to work out the equilibrium condition (lots of algebra involved) but feel free to predict what it is if you like.]
 - Is work done in the approach to equilibrium? Explain why or why not.
 - Does the internal energy U of the system change? Explain why or why not.



2. In class we considered the text example of a “polymer” comprising 4 monomer units that can exist in a “compact” state or an “open” state. The compact state arises from an attractive bond between the monomers at the ends of the chain (see figure). This bond is characterized by an energy $-\epsilon$. If this bond has an energy of 588×10^{-23} J, what is the free energy of the compact system at 310K? What is the free energy of the open system at 310K? Will there be more open or compact chains at 310K? The value of k is 1.38×10^{-23} J/K.
3. One gram of water is heated in a calorimeter from 274K to 314K. The following data are obtained:

Temperature	Cumulative Heat added (bomb calorimeter)	Total Heat added (constant pressure calorimeter)
274	0 cal	0 cal
294	19.7 cal	20 cal
314	39.5 cal	40 cal

What is the change in entropy when the temperature of 1 gm of water is raised from 274 to 314K at constant volume? At constant pressure?